



Jones, M. A. (2019) Retrograde signalling as an informant of circadian timing. *New Phytologist*, 221(4), pp. 1749-1753.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

This is the peer reviewed version of the following article Jones, M. A. (2019) Retrograde signalling as an informant of circadian timing. *New Phytologist*, 221(4), pp. 1749-1753, which has been published in final form at <http://dx.doi.org/10.1111/nph.15525>. This article may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

<http://eprints.gla.ac.uk/194057/>

Deposited on: 4 September 2019

1 Title: Retrograde Signalling as an Informant of Circadian Timing
2
3 Author: Matthew A. Jones
4
5 Address: School of Biological Sciences
6 University of Essex
7 Wivenhoe Park
8 Colchester
9 CO4 3SQ
10
11 Corresponding author: Dr Matt Jones
12 +44(0)1206 874740
13 matthew.jones@essex.ac.uk
14
15 Summary word count: 153
16 Total word count: 2115
17
18 Number of Figures: 1 (in colour)

19		
20	Summary	3
21	I) The circadian system is responsive to environmental change	4
22	II) Photoassimilates regulate circadian timing	5
23	III) Retrograde signals contribute to circadian timing.....	6
24	IV) Conclusions.....	9
25	V) Acknowledgements	9
26		
27		

28 **Summary**

29 The circadian system comprises of interlocking transcriptional/translational feedback loops
30 that regulate gene expression and consequently modulate plant development and physiology.
31 In order to maximize utility, the circadian system is entrained by changes in temperature and
32 light, allowing endogenous rhythms to be synchronized with both daily and seasonal
33 environmental change. While a great deal of environmental information is decoded by a suite
34 of photoreceptors, it is also becoming apparent that changes in cellular metabolism also
35 contribute to circadian timing, either through the stimulation of metabolic pathways or
36 through the accumulation of metabolic intermediates as a consequence of environmental
37 stress. As the source of many of these metabolic byproducts, mitochondria and chloroplasts
38 have begun to be viewed as environmental sensors, and rapid advancement of this field is
39 revealing the complex web of signalling pathways initiated by organelle perturbation. This
40 review will highlight recent advances in our understanding of how this metabolic regulation
41 influences circadian timing.

42 **Keywords**

43 Chloroplast

44 Circadian

45 MEcPP

46 PAP

47 Reactive Oxygen Species (ROS)

48 Retrograde

I) The circadian system is responsive to environmental change

Circadian timing modulates many biological processes, ranging from gene expression through to developmental decisions such as the transition to flowering (Millar, 2016). Although the circadian system is able to oscillate in the absence of environmental cues these endogenous rhythms are only useful if they are synchronized with and compensated against daily and seasonal changes in daylength and temperature (Millar, 2016). Plants perceive changes in their environment through the actions of photoreceptors and indicators of temperature, although this distinction is somewhat muddled as temperature and light signalling pathways share common components (Casal & Qüesta, 2018). Changes in temperature and light intensity (or quality) are sufficient to alter the expression of clock components, with the sensitivity of the circadian system to such environmental changes varying over the course of the day (Millar, 2016). This circadian ‘gating’ of responses to environmental change prevents the continuous resetting of the circadian timer to environmental inputs, while also allowing for the entrainment of the circadian clock to seasonal variation in daylength. Circadian gating also permits the modulation of plants’ responses to environmental change (for example by enabling greater response to chilling stress during the evening).

The nuclear circadian system comprises interlocked transcriptional/translational feedback loops that combine to generate oscillations of approximately 24 hrs (Millar, 2016). Successive waves of transcriptional activators and repressors regulate gene expression, with the Myb-like transcription factors CIRCADIAN CLOCK ASSOCIATED1 (CCA1) and LATE ELONGATED HYPOCOTYL (LHY) repressing expression at dawn whereas PSEUDORESPONSE REGULATOR9 (PRR9) and its orthologues PRR7/5/TOC1 limit expression during the day (Hsu & Harmer, 2014). Later in the evening, LUX ARRHYTHMO (LUX), EARLY FLOWERING3 (ELF3), and ELF4 associate to form an Evening Complex that similarly represses gene expression (Hsu & Harmer, 2014). Transcriptional activators including REVEILLE8 (RVE8), NIGHT LIGHT-INDUCIBLE (LNK) and LIGHT-REGULATED WD (LWD) proteins complement the activity of these transcriptional repressors (Hsu & Harmer, 2014). The combined activity of these feedback loops increases the amplitude of circadian rhythms and allows the clock to be compensated against seasonal changes in temperature, as well as improving the robustness of circadian rhythmicity (Shalit-Kaneh *et al.*, 2018).

While the transcriptional components of the circadian system are comparatively well understood, how these interconnecting feedback loops are influenced by environmental

change remains to be fully elucidated (Hsu & Harmer, 2014). The expression of dawn-phased clock components tends to be induced by light, while the Evening Complex has been identified as a likely hub for signal integration as it directly interacts with the photo- and thermo-sensor phytochromeB (phyB; (Hsu & Harmer, 2014; Huang, H *et al.*, 2016; Ezer *et al.*, 2017). The role of phytochrome, cryptochrome, and UVR8 photoreceptors in the circadian system has recently been discussed elsewhere (Oakenfull & Davis, 2017), and so this review will instead focus upon our understanding of how metabolic changes induced by environmental fluctuations affect circadian timing.

II) Photoassimilates regulate circadian timing

Plants' photosynthetic nature ensures that their metabolism is profoundly affected by the availability of light, with distinct changes in cellular energy levels and photoassimilates observed in the absence of this resource. Although the action of photoreceptors is sufficient to entrain and maintain circadian rhythms (Millar, 2016) the metabolic consequences of photosynthesis also contribute to circadian rhythms. Recent work has documented how the accumulation of sucrose (and other photoassimilates) is sufficient to reset the circadian system (Dalchau *et al.*, 2011; Haydon *et al.*, 2013; Haydon *et al.*, 2017). Products from photosynthesis are integrated with the circadian oscillator via at least two pathways, with one acting to alter phase in a PRR7- and CCA1-dependent manner (Haydon *et al.*, 2013), whereas GIGANTEA promotes circadian rhythmicity in constant darkness in the presence of sucrose (Dalchau *et al.*, 2011; Haydon *et al.*, 2017).

One possible mechanism linking photoassimilates with the circadian system revolves around the SnRK1 signalling hub (Fig. 1). SnRK1 regulates metabolism by phosphorylating metabolic enzymes, with its activity regulated in part through control of the AKIN10 catalytic subunit (Wurzinger *et al.*, 2018). SnRK1 activity is additionally repressed by the accumulation of trehalose-6-phosphate, which serves as a molecular indicator of intracellular sucrose (Figueroa & Lunn, 2016). Over-expression of AKIN10 extends circadian period in the light (Shin *et al.*, 2017; Frank *et al.*, 2018), suggesting that control of this subunit is sufficient to change circadian timing. Importantly, SnRK1 regulates the activity of bZIP63, a transcription factor that activates *PRR7* transcription (Mair *et al.*, 2015; Frank *et al.*, 2018). As a consequence, regulation of SnRK1 activity by trehalose-6-phosphate provides a mechanism by which intracellular sugar levels can be interpreted by the circadian system (Frank *et al.*, 2018).

III) Retrograde signals contribute to circadian timing

Cell function is ultimately a cooperation between distinct organelles, which is maintained by both anterograde and retrograde signalling pathways (recently comprehensively reviewed by (Chan *et al.*, 2016b; de Souza *et al.*, 2017). Retrograde signals convey information provided by organelles such as chloroplasts and mitochondria to the nucleus, ultimately altering nuclear gene expression. For example, mature chloroplasts contribute to lipid, starch, sulfur, and amino acid metabolism, as well as contributing to the production of several hormones, in addition to their primary role in photosynthesis (Chan *et al.*, 2016b; de Souza *et al.*, 2017). Similarly, mitochondria are essential for cellular respiration and contribute to the generation of reactive oxygen species such as $^1\text{O}_2$ and H_2O_2 (Wang *et al.*, 2018). Retrograde signals consequently relay the metabolic health of the cell to the nucleus, and so provide information useful for integration into the circadian oscillator in addition to the signalling provided by photo- and thermo-sensors.

During chloroplast biogenesis, several pathways relay the general health and developmental status of the plastid to the nucleus, resulting in changes in nuclear gene expression as a consequence of biogenic control (Chan *et al.*, 2016b; de Souza *et al.*, 2017). Disruption of chloroplast function is sufficient to alter photosynthesis-associated nuclear genes (PhANGs) and nuclear circadian gene expression. For instance, loss of CHLOROPLAST RNA BINDING (CRB) delays phase and increases the amplitude of circadian gene expression (Hassidim *et al.*, 2007), while application of norflurazon (an inhibitor of photosynthesis) and lincomycin (which inhibits plastid protein synthesis) extends circadian period (Chen *et al.*, 2013). Similarly, an iron deficiency extends circadian period likely due to gross disruption of chloroplast function (Chen *et al.*, 2013; Salomé *et al.*, 2013). It is therefore apparent that retrograde signals relaying the health of the chloroplast are sufficient to delay circadian timing.

In addition to signals relaying the developmental status of organelles, the operation of mature chloroplasts can be perturbed by environmental factors (Fig. 1). Increases in irradiance promote photosynthesis and induce nonphotochemical quenching whereas changing temperatures govern the speed of enzymatic reactions. Such environmental stressors are often sufficient to induce metabolic imbalances, causing oxidative stress in mitochondria and chloroplasts and leading to changes to the energy status of the cell (Chan *et al.*, 2016b). Such disruption of metabolism additionally perturbs hormone biosynthesis and signalling, alters calcium signalling, and induces the accumulation of metabolic intermediaries that can subsequently be transported from the mitochondria or chloroplasts to

the nucleus (Chan *et al.*, 2016b; Mullineaux *et al.*, 2018). With regards to circadian rhythmicity, high temperatures are sufficient to slow the circadian oscillator, while the circadian system is also delayed by osmotic stress (Gil *et al.*, 2017; Litthauer *et al.*, 2018). As with many stress responses, the question now facing the field is whether the clock is slowed under these environmental conditions as part of a generalized reaction to cellular damage, or whether specific retrograde signals delay circadian progression as part of an adaptive response.

Reactive Oxygen Species

Reactive oxygen species (ROS) arise from the metabolic processes of mitochondria and chloroplasts (Murchie & Niyogi, 2011; Huang, S *et al.*, 2016). Both H₂O₂ accumulation and the oxidation state of peroxiredoxins (a ubiquitous family of enzymes that serve as antioxidants and regulatory proteins) vary with a circadian rhythm, suggesting that ROS generation is influenced by the circadian system (Edgar *et al.*, 2012; Lai *et al.*, 2012). Although ROS are produced as a consequence of the normal functioning of photosynthesis and respiration their accumulation increases during sub-optimal conditions, when environmental stresses such as high light induce an imbalance between photosynthetic and respiratory pathways (Mullineaux *et al.*, 2018). Under these stressful conditions, an increase in ROS production causes significant damage to the metabolic machinery, inducing photoinhibition and a generalized damage response (Mullineaux *et al.*, 2018). ROS such as H₂O₂ can be transferred directly to the nucleus from the chloroplast (Exposito-Rodriguez *et al.*, 2017), where it is likely that the activity of redox-sensitive transcription factors is modulated. However, it remains unknown how these factors contribute to circadian timing. By contrast, shorter lived ROS such as ¹O₂ are likely to initiate signalling pathways by oxidizing carotenoids or polyunsaturated fatty acids (Mullineaux *et al.*, 2018). One such signalling trigger is β-Cyclocitral, which is an oxidised derivative of β-carotene that accumulates as a consequence of ¹O₂ production and which induces gene expression in response to high light (Ramel *et al.*, 2012).

In addition to the direct role of ROS, or oxidized by-products, plants also have a suite of mechanisms, including non-photochemical quenching, that allow for the sequestration of ROS below critical thresholds (Ruban, 2016). Above these thresholds, ROS accumulation leads to the production of a range of metabolites that plants subsequently utilize as signalling molecules to report the metabolic status of the mitochondria and chloroplast. Such signals could be integrated into the nuclear circadian oscillator and include molecules such as

methylethanol cyclodiphosphate (MEcPP), and 5'-Phosphoadenosine 3'-phosphate (PAP). Although these metabolites are now becoming accepted as authentic retrograde signals, it is not yet clear how these molecules integrate with circadian timing (Fig. 1).

MEcPP

Methylethanol cyclodiphosphate (MEcPP) accumulates in response to wounding or high light stress, inducing the expression of nuclear stress-responsive genes and repressing the accumulation of auxin (Xiao *et al.*, 2012; Jiang *et al.*, 2018). Our understanding of MEcPP is still developing, but one exemplar signalling pathway repressed by MEcPP accumulation involves the transcription factor BBX19. *BBX19* transcription is repressed in *cehl* mutants that constitutively accumulate MEcPP as a consequence of disruption of the methylethanol phosphate pathway within the plastid (Xiao *et al.*, 2012; Wang *et al.*, 2014). BBX19 has an important role in photomorphogenesis, repressing *PIF4* and *PIF5* expression by promoting the turnover of ELF3 (Wang *et al.*, 2015). As a consequence, MEcPP accumulation results in the accumulation of ELF3. Since ELF3-overexpressing lines have previously been reported as having a long circadian period (Covington *et al.*, 2001) it is plausible that MEcPP accumulation could result in extended circadian rhythms, although this hypothesis has yet to be tested (Fig. 1).

PAP

3'-PhosphoAdenosine 5'-Phosphate (PAP) is a by-product of sulfur metabolism that is usually metabolised by SAL1, a redox-sensitive phosphatase that accumulates in both chloroplasts and mitochondria (Estavillo *et al.*, 2011; Chan *et al.*, 2016a). Following the application of high light or drought stress, SAL1 becomes oxidized, leading to its inactivation and the consequential accumulation of PAP (Estavillo *et al.*, 2011; Chan *et al.*, 2016a). PAP accumulation has several metabolic consequences, including the inactivation of XRN ribonucleases (Dichtl *et al.*, 1997; Mechold *et al.*, 2006; Fig. 1). Loss of XRN function leads to the accumulation of uncapped RNAs such as those generated from miRNA processing (Kurihara *et al.*, 2012). Interestingly, either the accumulation of PAP or the loss of XRN activity is sufficient to extend circadian period (Litthauer *et al.*, 2018). Although the mechanism underlying this phenotype remains unclear, such data are consistent with previous reports that gross defects in RNA processing slow circadian progression (Nolte & Staiger, 2015). It will therefore be important to examine how loss of XRN activity alters the stability

of circadian transcripts, and to understand how such post-transcriptional regulation acts in concert with previously reported variations in splicing to control functional gene expression.

IV) Conclusions

Plants are highly sensitive to environmental change, and we have identified a complex suite of photo- and thermo-sensors that enable these responses, as well as the circadian system that anticipates and integrates these signals to regulate development and plants responses. However, it is also important to note the contribution of metabolism-induced signalling pathways to plants responses. We are beginning to appreciate how photosynthesis and metabolic perturbations within organelles lead to differential nuclear gene expression, and consequently to circadian timing. Understanding how mitochondrial and chloroplastic signalling contribute to the circadian system (and how these pathways are reciprocally modulated by the circadian system) will allow the development of a holistic understanding of plants' responses to environmental change.

V) Acknowledgements

Research in the Jones lab is supported by the University of Essex. The author apologies to colleagues whose work could not be directly cited due to space limitations.

VI) References

- Casal JJ, Qüesta JI. 2018. Light and temperature cues: multitasking receptors and transcriptional integrators. *New Phytologist* **217**(3): 1029-1034.
- Chan KX, Mabbitt PD, Phua SY, Mueller JW, Nisar N, Gigolashvili T, Stroeher E, Grassl J, Arlt W, Estavillo GM, et al. 2016a. Sensing and signaling of oxidative stress in chloroplasts by inactivation of the SAL1 phosphoadenosine phosphatase. *PNAS* **113**(31): E4567-4576.
- Chan KX, Phua SY, Crisp P, McQuinn R, Pogson BJ. 2016b. Learning the Languages of the Chloroplast: Retrograde Signaling and Beyond. *Annual Review of Plant Biology* **67**: 25-53.
- Chen Y-Y, Wang Y, Shin L-J, Wu J-F, Shanmugam V, Tsednee M, Lo J-C, Chen C-C, Wu S-H, Yeh K-C. 2013. Iron is involved in the maintenance of circadian period length in Arabidopsis. *Plant Physiology* **161**(3): 1409-1420.
- Covington MF, Panda S, Liu XL, Strayer CA, Wagner DR, Kay SA. 2001. ELF3 modulates resetting of the circadian clock in Arabidopsis. *Plant Cell* **13**(6): 1305-1315.
- Dalchau N, Baek SJ, Briggs HM, Robertson FC, Dodd AN, Gardner MJ, Stancombe MA, Haydon MJ, Stan G-B, Gonçalves JM, et al. 2011. The circadian oscillator gene *GIGANTEA* mediates a long-term response of the Arabidopsis thaliana circadian clock to sucrose. *PNAS* **108**(12): 5104-5109.

- de Souza A, Wang J-Z, Dehesh K. 2017.** Retrograde signals: Integrators of interorganellar communication and orchestrators of plant development. *Annual Review of Plant Biology* **68**: 85-108.
- Dichtl B, Stevens A, Tollervey D. 1997.** Lithium toxicity in yeast is due to the inhibition of RNA processing enzymes. *EMBO Journal* **16**(23): 7184-7195.
- Edgar RS, Green EW, Zhao Y, Van Ooijen G, Olmedo M, Qin X, Xu Y, Pan M, Valekunja UK, Feeney KA, et al. 2012.** Peroxiredoxins are conserved markers of circadian rhythms. *Nature* **485**(7399): 459-464.
- Estavillo GM, Crisp PA, Pornsiriwong W, Wirtz M, Collinge D, Carrie C, Giraud E, Whelan J, David P, Javot H, et al. 2011.** Evidence for a SAL1-PAP chloroplast retrograde pathway that functions in drought and high light signaling in Arabidopsis. *Plant Cell* **23**(11): 3992-4012.
- Exposito-Rodriguez M, Laissue PP, Yvon-Durocher G, Smirnov N, Mullineaux PM. 2017.** Photosynthesis-dependent H₂O₂ transfer from chloroplasts to nuclei provides a high-light signalling mechanism. *Nature Communications* **8**(1): 49.
- Ezer D, Jung J-H, Lan H, Biswas S, Gregoire L, Box MS, Charoensawan V, Cortijo S, Lai X, Stöckle D, et al. 2017.** The evening complex coordinates environmental and endogenous signals in Arabidopsis. *Nature Plants* **3**: 17087.
- Figueroa CM, Lunn JE. 2016.** A Tale of Two Sugars: Trehalose 6-Phosphate and Sucrose. *Plant Physiology* **172**(1): 7-27.
- Frank A, Mantioli CC, Webb AAR, Dodd AN. 2018.** Circadian entrainment in Arabidopsis by the sugar-responsive transcription factor bZIP63. *Current Biology* **28**(16): 2597-2606.
- Gil K-E, Kim W-Y, Lee H-J, Faisal M, Saquib Q, Alatar AA, Park C-M. 2017.** ZEITLUPE contributes to a thermoresponsive protein quality control system in Arabidopsis. *Plant Cell* **29**(11): 2882-2894.
- Hassidim M, Yakir E, Fradkin D, Hilman D, Kron I, Keren N, Harir Y, Yerushalmi S, Green R. 2007.** Mutations in *CHLOROPLAST RNA BINDING* provide evidence for the involvement of the chloroplast in the regulation of the circadian clock in Arabidopsis. *Plant J* **51**(4): 551-562.
- Haydon MJ, Mielczarek O, Frank A, Román Á, Webb AA. 2017.** Sucrose and ethylene signaling interact to modulate the circadian clock. *Plant Physiology* **175**(2): 947-958.
- Haydon MJ, Mielczarek O, Robertson FC, Hubbard KE, Webb AAR. 2013.** Photosynthetic entrainment of the Arabidopsis thaliana circadian clock. *Nature* **502**(7473): 689-692.
- Hsu PY, Harmer SL. 2014.** Wheels within wheels: The plant circadian system. *Trends in Plant Science* **19**(4): 240-249.
- Huang H, Alvarez S, Bindbeutel R, Shen Z, Naldrett MJ, Evans BS, Briggs SP, Hicks LM, Kay SA, Nusinow DA. 2016.** Identification of Evening Complex-associated proteins in Arabidopsis by affinity purification and mass spectrometry. *Molecular & Cellular Proteomics* **15**(1): 201-217.
- Huang S, Van Aken O, Schwarzländer M, Belt K, Millar AH. 2016.** The Roles of Mitochondrial reactive oxygen species in cellular signaling and stress response in plants. *Plant Physiology* **171**(3): 1551-1559.
- Jiang J, Rodriguez-Furlan C, Wang J-Z, de Souza A, Ke H, Pasternak T, Lasok H, Ditengou FA, Palme K, Dehesh K. 2018.** Interplay of the two ancient metabolites auxin and MEcPP regulates adaptive growth. *Nature Communications* **9**(1): 2262.

- Kurihara Y, Schmitz RJ, Nery JR, Schultz MD, Okubo-Kurihara E, Morosawa T, Tanaka M, Toyoda T, Seki M, Ecker JR. 2012. Surveillance of 3' Noncoding Transcripts Requires FIERY1 and XRN3 in Arabidopsis. *G3 (Bethesda, Md.)* 2(4): 487-498.
- Lai AG, Doherty CJ, Mueller-Roeber B, Kay SA, Schippers JHM, Dijkwel PP. 2012. CIRCADIAN CLOCK-ASSOCIATED 1 regulates ROS homeostasis and oxidative stress responses. *PNAS* 109(42): 17129-17134.
- Litthauer S, Chan KX, Jones MA. 2018. 3'-Phosphoadenosine 5'-phosphate accumulation delays the circadian system. *Plant Physiology* 176(4): 3120-3135.
- Mair A, Pedrotti L, Wurzinger B, Anrather D, Simeunovic A, Weiste C, Valerio C, Dietrich K, Kirchler T, Nägele T, et al. 2015. SnRK1-triggered switch of bZIP63 dimerization mediates the low-energy response in plants. *eLife* 4: e05828
- Mechold U, Ogryzko V, Ngo S, Danchin A. 2006. Oligoribonuclease is a common downstream target of lithium-induced pAp accumulation in Escherichia coli and human cells. *Nucleic Acids Research* 34(8): 2364-2373.
- Millar AJ. 2016. The intracellular dynamics of circadian clocks reach for the light of ecology and evolution. *Annual Review of Plant Biology* 67: 595-618.
- Mullineaux PM, Exposito-Rodriguez M, Laissue PP, Smirnoff N. 2018. ROS-dependent signalling pathways in plants and algae exposed to high light: Comparisons with other eukaryotes. *Free Radical Biology & Medicine* 122: 52-64.
- Murchie EH, Niyogi KK. 2011. Manipulation of photoprotection to improve plant photosynthesis. *Plant Physiology* 155(1): 86-92.
- Nolte C, Staiger D. 2015. RNA around the clock - regulation at the RNA level in biological timing. *Frontiers in Plant Science* 6: 311.
- Oakenfull RJ, Davis SJ. 2017. Shining a light on the Arabidopsis circadian clock. *Plant, Cell & Environment* 40(11): 2571-2585.
- Ramel F, Birtic S, Ginies C, Soubigou-Taconnat L, Triantaphylidès C, Havaux M. 2012. Carotenoid oxidation products are stress signals that mediate gene responses to singlet oxygen in plants. *PNAS* 109(14): 5535-5540.
- Ruban AV. 2016. Nonphotochemical chlorophyll fluorescence quenching: Mechanism and effectiveness in protecting plants from photodamage. *Plant Physiology* 170(4): 1903-1916.
- Salomé PA, Oliva M, Weigel D, Krämer U. 2013. Circadian clock adjustment to plant iron status depends on chloroplast and phytochrome function. *EMBO Journal* 32(4): 511-523.
- Shalit-Kaneh A, Kumimoto RW, Filkov V, Harmer SL. 2018. Multiple feedback loops of the Arabidopsis circadian clock provide rhythmic robustness across environmental conditions. *PNAS* 115(27): 7147-7152.
- Shin J, Sanchez-Villarreal A, Davis AM, Du S-X, Berendzen KW, Koncz C, Ding Z, Li C, Davis SJ. 2017. The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent manner. *Plant, Cell & Environment* 40(7): 997-1008.
- Wang C-Q, Guthrie C, Sarmast MK, Dehesh K. 2014. BBX19 interacts with CONSTANS to repress *FLOWERING LOCUS T* transcription, defining a flowering time checkpoint in Arabidopsis. *Plant Cell* 26(9): 3589-3602.
- Wang C-Q, Sarmast MK, Jiang J, Dehesh K. 2015. The transcriptional regulator BBX19 promotes hypocotyl growth by facilitating COP1-mediated EARLY FLOWERING3 degradation in Arabidopsis. *Plant Cell* 27(4): 1128-1139.

- Wang Y, Berkowitz O, Selinski J, Xu Y, Hartmann A, Whelan J. 2018. Stress responsive mitochondrial proteins in *Arabidopsis thaliana*. *Free Radical Biology & Medicine* **122**: 28-39.
- Wurzinger B, Nukarinen E, Nägele T, Weckwerth W, Teige M. 2018. The SnRK1 kinase as central mediator of energy signaling between different organelles. *Plant Physiology* **176**(2): 1085-1094.
- Xiao Y, Savchenko T, Baidoo EEK, Chehab WE, Hayden DM, Tolstikov V, Corwin JA, Kliebenstein DJ, Keasling JD, Dehesh K. 2012. Retrograde signaling by the plastidial metabolite MEcPP regulates expression of nuclear stress-response genes. *Cell* **149**(7): 1525-1535.

Figure Legends

Figure 1. Operational retrograde signalling contributes to circadian timing in response to environmental factors. The nuclear circadian oscillator consists of multiple negative feedback loops (including Myb transcription factors, PRRs, and the Evening Complex) in complex with activating proteins (summarized in Hsu & Harmer, 2014; Millar, 2016). These transcriptional/translational feedback loops are modified by metabolic intermediaries and products. **(a)** The sucrose/Tre6P nexus regulates the activity of SnRK1 dependent on cytosolic sucrose accumulation. The SnRK1 complex phosphorylates bZIP63, allowing this transcription factor to dimerize and promote the expression of *PRR7*. **(b, c)** The infliction of abiotic stress induces the accumulation of ROS that alters the chloroplast redox state such that MEcPP and PAP accumulate. MEcPP represses the expression of *BBX19*, leading to the accumulation of ELF3 which is a core constituent of the Evening Complex. PAP accumulation inhibits XRN activity, leading to defects in mRNA processing and an extension of circadian period. Abbreviations: ELF3, EARLY FLOWERING3; MEcPP, methylerythritol cyclodiphosphate; PAP, 5'-PhosphoAdenosine 3'-Phosphate; PRRs, PSEUDORESPONSE REGULATORS; ROS, Reactive Oxygen Species; Tre6P, trehalose 6-phosphate; XRN, EXORIBONUCLEASE.

